

APPROXIMATE CONNES-AMENABILITY OF DUAL BANACH ALGEBRAS

G. H. Esslamzadeh and B. Shojaei

Abstract. We introduce the notions of approximate Connes-amenability and approximate strong Connes-amenability. Then we characterize these two types of dual Banach algebras in terms of approximate normal virtual diagonal and approximate σWC -virtual diagonals. Some concrete cases are also discussed.

Mathematics Subject Classification (2000). Primary 46H25, 46H20; Secondary 46H35.

Keywords. Approximately inner derivation, Approximately weakly amenable, Approximately amenable, Approximate trace extension property.

1. INTRODUCTION

In [9], B.E. Johnson, R. V. Kadison, and J. Ringrose introduced a notion of amenability for von Neumann algebras which modified Johnson's original definition for Banach algebras [8] in the sense that it takes the dual space structure of a von Neumann algebra into account. This notion of amenability was later called Connes-amenability by A. Ya. Helemskii [7]. In [10], the author extended the notion of Connes-amenability to the larger class of dual Banach algebras. The concept of approximate amenability for a Banach algebras was introduced by F. Ghahramani and R. J. Loy in [5]. Our motivation for introducing approximate Connes-amenability is finding a version of these approximate forms of amenability which sounds suitable for dual Banach algebras.

Before proceeding further we recall some terminology.

The main part of this work was undertaken while the second author was studying for PhD at the Islamic Azad University. The second author wishes to thank Islamic Azad University for the financial support.

Throughout A is a Banach algebra and X is a Banach A -bimodule. A derivation is a bounded linear map $D : A \longrightarrow X$ such that

$$D(ab) = a.Db + Da.b \quad (a, b \in A).$$

For $x \in X$, define

$$\delta_x(a) = a.x - x.a \quad (a \in A).$$

Then δ_x is a derivation; maps of this form are called inner derivation. A derivation $D : A \longrightarrow X$ is approximately inner if there exists a net $(a_\alpha) \subseteq X$ such that for every $a \in A$, $D(a) = \lim_\alpha (a.x_\alpha - x_\alpha.a)$, the limit being in norm. We say that A is approximately amenable if for any A -bimodule X , every derivation $D : A \longrightarrow X^*$ is approximately inner. A is called a dual Banach algebra if there is a closed submodule A_* of A^* such that $A = (A_*)^*$. For example if G is a locally compact group, then $M(G)$ is a dual Banach algebra (with $A_* = c_0(G)$), or if A is an Arens regular Banach algebra then A^{**} is a dual Banach algebra.

Let A be a dual Banach algebra. A dual Banach A -bimodule X is called normal if, for each $x \in X$, the maps

$$A \longrightarrow X, \quad a \longmapsto \begin{cases} a.x \\ x.a \end{cases}$$

are ω^* -continuous. A dual Banach algebra A is Connes-amenable if, for every normal, dual Banach A -module X , every ω^* -continuous derivation $D : A \longrightarrow X$ is inner.

Let A be a dual Banach algebra, and let X be a Banach A -bimodule. Then we call an element $\phi \in X^*$ a ω^* -element if the maps

$$A \longrightarrow X^*, \quad a \longmapsto \begin{cases} a.\phi \\ \phi.a \end{cases}$$

are $\omega^* - \omega^*$ continuous.

2. APPROXIMATE CONNES AMENABILITY

Definition 2.1. A dual Banach algebra A is approximately Connes-amenable if for every normal, dual Banach A -bimodule X , every ω^* -continuous derivation $D \in Z^1(A, X)$ is approximately inner.

Lemma 2.2. Suppose that A is approximately Connes-amenable. Then A has left and right approximate identities. In particular A^2 is dense in A .

Proof. Let X be the Banach A -bimodule whose underlying linear space is A equipped with the following module operation:

$$a.x = ax \quad \text{and} \quad x.a = 0 \quad (a \in A, \quad x \in X).$$

Obviously X , is a normal dual Banach A -bimodule and the identity map on A is a ω^* -continuous derivation. Since A is approximately Connes-amenable, then there exists a net $(a_\alpha) \subseteq X$ such that

$$a = \lim_{\alpha} aa_{\alpha} \quad (a \in A).$$

This means that A has a right approximate identity. Similarly, one see that A has a left approximate identity. $\square$

Let (A, A_*) be a dual Banach algebra, and let $A^\#$ be the Banach algebra $A \oplus C$. Then $A^\#$ is a dual Banach algebra with predual $A_* \oplus C$ and norm

$$\|(\mu, \alpha)\| = \max(\|\mu\|, |\alpha|) \quad (\mu \in A_*, \alpha \in C).$$

Proposition 2.3. Let A be a dual Banach algebra. A is approximately Connes-amenable if and only if $A^\#$ is approximately Connes-amenable.

Proof. Let $D : A^\# \longrightarrow X^*$ be a ω^* -continuous derivation for normal dual Banach $A^\#$ -bimodule X^* . By [5, Lemma 2.3], $D = D_1 + ad_\eta$ where $D_1 : A^\# \longrightarrow e.X^*.e$ is a ω^* -continuous derivation and $\eta \in X^*$. Since $e.X^*.e$ is a normal dual Banach A -bimodule, then $D_1(e) = 0$ and $D_1|_A$ is approximately inner; whence D is approximately inner. Thus $A^\#$ is approximately Connes-amenable.

Now suppose $D : A \longrightarrow X^*$ is a ω^* -continuous derivation for normal dual Banach A -bimodule X^* . Set

$$\tilde{D} : A^\# \longrightarrow X^*, \quad \tilde{D}(a + \lambda e) = Da \quad (a \in A, \lambda \in C).$$

If we define $e.x = x.e = x$ ($e \in A^\#, x \in X$), then X^* turns into a normal dual Banach $A^\#$ -bimodule and $\tilde{D}$ is a ω^* -continuous derivation. So $\tilde{D}$ is approximately inner, and hence so is D . It follows that A is approximately Connes-amenable. $\square$

Proposition 2.4. Let A be an Arens regular dual Banach algebra. If A^{**} is approximately Connes-amenable, then A is approximately Connes-amenable.

Proof. Suppose X is a normal dual Banach A -bimodule, and $\pi : A^{**} \longrightarrow A$ is the restriction map to A_* . Then π is a $\omega^* - \omega^*$ continuous homomorphism. Therefore X is a normal dual Banach A^{**} -bimodule with the following actions

$$a^{**}.x = \pi(a^{**})x \quad , \quad x.a^{**} = x\pi(a^{**}) \quad (x \in X, a^{**} \in A^{**}).$$

Let $D : A \longrightarrow X$ be a $\omega^* - \omega^*$ continuous derivation. It is easy to see that $Do\pi : A^{**} \longrightarrow X$ is a $\omega^* - \omega^*$ continuous derivation. Since A^{**} is approximately Connes-amenable, then there exists a net $(x_\alpha) \subseteq X$ such that

$$Do\pi(a^{**}) = \lim_{\alpha} a^{**}.x_\alpha - x_\alpha.a^{**} \quad (a^{**} \in A^{**}).$$

So

$$D(a) = \lim_{\alpha} a.x_\alpha - x_\alpha.a \quad (a \in A).$$

□

For a locally compact group G we have:

Theorem 2.5. If $L^1(G)$ is approximately amenable then $M(G)$ is approximately Connes-amenable.

Proof. By [5, Theorem 3.2], G is amenable, and by [11, Theorem 4.4.13], $M(G)$ is approximately Connes-amenable. □

Theorem 2.6. Let A be an Arens regular Banach algebra which is an ideal in A^{**} . If A^{**} is approximately Connes-amenable and has an identity then A is approximately amenable.

Proof. By [5, Proposition 2.5] in order to show that A is approximately amenable it is sufficient to show that every $D \in Z^1(A, X^*)$ is approximately inner for each neo-unital Banach A -module.

Let X be a neo-unital Banach A -bimodule, and let $D \in Z^1(A, X^*)$. By [11, Theorem 4.4.8] X^* is a normal dual Banach A^{**} -bimodule and D has a unique extension $\tilde{D} \in Z^1(A^{**}, X^*)$. From the approximate Connes-amenability of A^{**} we conclude that $\tilde{D}$, and hence D is inner. It follows that A is approximately amenable. □

In the following proposition we obtain a criterion for approximate amenability which will be used in the sequel.

Proposition 2.7. The following conditions are equivalent;

- (i) A is approximately amenable;
- (ii) For any A -bimodule X , every bounded derivation $D : A \longrightarrow X^{**}$ is approximately inner.

Proof (i) $\implies$ (ii) This is immediate.

(ii) $\implies$ (i). By [6, Theorem 2.1], it suffices to show that if $D \in Z^1(A, X)$, then it is approximately inner.

We have $\iota \circ D \in Z^1(A, X^{**})$, where $\iota : X \longrightarrow X^{**}$ is the canonical embedding.

Thus, there exists a net $(x_\alpha^{**}) \subseteq X^{**}$ such that

$$\iota \circ D(a) = \lim_{\alpha} a.x_\alpha^{**} - x_\alpha^{**}.a \quad (a \in A).$$

Now take $\epsilon > 0$, and finite sets $F \subseteq A$, $\phi \subseteq X^*$.

Then there is α such that

$$|\langle x^*, \iota \circ D(a) - (a.x_\alpha^{**} - x_\alpha^{**}.a) \rangle| < \epsilon$$

For all $x^* \in \phi$ and $a \in F$.

By Goldstine's Theorem, there is a net (x_α) in X such that

$$|\langle x^*, \iota \circ D(a) - (a.x_\alpha - x_\alpha.a) \rangle| < \epsilon$$

for all $x^* \in \phi$ and $a \in F$. Thus there is a net $(x_\alpha) \subseteq X$ such that

$$Da = \omega - \lim_{\alpha} a.x_\alpha - x_\alpha.a \quad (a \in A).$$

Finally, for each finite set $F \subseteq A$, say $F = \{a_1, \dots, a_n\}$,

$$(a_1.x_\alpha - x_\alpha.a_1, \dots, a_n.x_\alpha - x_\alpha.a_n) \longrightarrow (Da_1, \dots, Da_n)$$

weakly in $(X)^n$. By Mazure's Theorem,

$$(Da_1, \dots, Da_n) \in \overline{CO}^{\|\cdot\|} \{ (a_1.x_\alpha - x_\alpha.a_1, \dots, a_n.x_\alpha - x_\alpha.a_n) \},$$

Thus there is a convex linear combination $x_{F,\epsilon}$ of elements in the set $\{x_\alpha\}$ such that ,

$$\|Da - (a.x_{F,\epsilon} - x_{F,\epsilon}.a)\| < \epsilon \quad (a \in F).$$

The family of such pairs (F, ϵ) is a directed set for the partial order $\leq$ given by

$$(F_1, \epsilon_1) \leq (F_2, \epsilon_2) \quad \text{if} \quad F_1 \subseteq F_2 \quad \text{and} \quad \epsilon_1 \geq \epsilon_2.$$

Thus

$$Da = \lim_{(F, \epsilon)} a.x_{(F, \epsilon)} - x_{(F, \epsilon)}.a \quad (a \in A).$$

□

One might ask whether the ideal condition in the Theorem 2.6 is necessary for A.

We answer this question, partially.

Theorem 2.8. Let A be an Arens regular Banach algebra which is a right ideal in A^{**} . Let for every A^{**} -bimodule X, $X^*A = X^*$. If A^{**} is approximately Connes-amenable, then A is approximately amenable.

Proof. Let X be a Banach A-bimodule. By proposition 2.7 it suffices to show that every $D \in Z^1(A, X^{**})$ is approximately inner.

Let $D \in Z^1(A, X^{**})$. By [2, p. 27], X^{****} is a Banach A^{**} -bimodule and by [1, Proposition 2.7.17(i)],

$$D^{**} : A^{**} \longrightarrow X^{****}$$

is a $\omega^* - \omega^*$ continuous derivation. Since X^{****} is an A^{**} -bimodule, then by assumption

$$X^{****}A = X^{****}. \quad (1)$$

We claim that X^{****} is a normal dual Banach A^{**} -bimodule. Let (a''_α) be a net in A^{**} such that $a''_\alpha \xrightarrow{\omega^*} a''$. Then

$$aa''_\alpha \xrightarrow{\omega^*} aa'' \quad (a \in A).$$

Since A is a right ideal of A^{**} and ω^* -topology of A^{**} restricted to A coincides with the weak topology, we have

$$aa''_\alpha \xrightarrow{\omega} aa'' \quad (a \in A).$$

Let $x'''' \in X^{****}$. Then by (1) there exists $a \in A$ and $y'''' \in X^{****}$ such that $x'''' = y''''a$. Thus we have,

$$x''''a''_\alpha = y''''a''_\alpha \xrightarrow{\omega} y''''aa'' = x''''aa''.$$

Therefore

$$x''''a''_\alpha \xrightarrow{\omega^*} x''''a''.$$

On the other hand by definition,

$$a''_{\alpha} x'''' \longrightarrow^{\omega^*} a'' x''''$$

Now since A^{**} is approximately Connes-amenable, then there exists a net $(x''''_{\alpha}) \in X''''$, such that

$$D^{**} a'' = \lim_{\alpha} a'' . x''''_{\alpha} - x''''_{\alpha} . a'' \quad (a'' \in A^{**}).$$

Let $P : X'''' \longrightarrow X''$ be the natural projection, then

$$Da = \lim_{\alpha} a . P(x''''_{\alpha}) - P(x''''_{\alpha}) . a \quad (a \in A).$$

Thus D is approximately inner. It follow that A is approximately amenable. $\square$

Proposition 2.9. Suppose that A is a dual Banach algebra with identity. Then A is approximately Connes-amenable if and only if every ω^* -continuous derivation into every unital normal dual Banach bimodule X is approximately inner.

Proof. ($\implies$): Obvious.

($\impliedby$) Suppose $D \in Z^1(A, X)$ is a $\omega^* - \omega^*$ continuous derivation into the normal dual Banach bimodule X . By [5, Lemma 2.3], we have $D = D_1 + ad_{\eta}$ where $D_1 : A \longrightarrow e.X.e$ is a derivation and $\eta \in X^*$. Since D is a ω^* -continuous derivation and X is a normal dual Banach bimodule then D_1 is ω^* -continuous and $e.X.e$ is normal. So by assumption D_1 is approximately inner, and therefore A is approximately Connes-amenable. $\square$

Let A be a dual Banach algebra with identity, and let $L^2_{\omega^*}(A, C)$ be the space of separately $\omega^* - \omega^*$ -continuous bilinear functionals on A . Clearly, $L^2_{\omega^*}(A, C)$ is a Banach A -submodule of $L^2(A, C)$ and

$$L^2(A, C) \simeq (A \widehat{\otimes} A)^*.$$

From [10] we have a natural A -bimodule map

$$\theta : A \otimes A \longrightarrow L^2_{\omega^*}(A, C)^*$$

defined by letting $\theta(a \otimes b)F = F(a, b)$. Since $A_* \otimes A_* \subseteq L^2_{\omega^*}(A, C)$ and $A_* \otimes A_*$ separates points of $A \otimes A$, then θ is one-to-one. We will identify $A \otimes A$ with its image, writing

$$A \otimes A \subseteq L^2_{\omega^*}(A, C)^*.$$

The map Δ_A is defined as follow:

$$\Delta_A : A \widehat{\otimes} A \longrightarrow A, \quad a \otimes b \longmapsto ab \quad (a, b \in A)$$

Since multiplication in a dual Banach algebra is separately $\omega^* - \omega^*$ -continuous, we have,

$$\Delta_A^* A_* \subset L_{\omega^*}^2(A, C),$$

So that restriction, of Δ_A^{**} to $L_{\omega^*}^2(A, C)$ turns into a Banach A -bimodule homomorphism

$$\Delta_{\omega^*} : L_{\omega^*}^2(A, C)^* \longrightarrow A$$

Suppose $F \in L_{\omega^*}^2(A, C)$ and $M \in L_{\omega^*}^2(A, C)^*$. We have the notation,

$$\int F(a, b) dM(a, b) = \int F dM := \langle M, F \rangle.$$

More generally given a dual Banach space X^* and a bounded bilinear function $F : A \times A \longrightarrow X^*$ such that $a \longrightarrow F(a, b)$ and $b \longrightarrow F(a, b)$ are $\omega^* - \omega^*$ -continuous, we define $\int F dM \in X^*$ by

$$\langle \int F dM, x \rangle = \int \langle F(a, b), x \rangle dM(a, b) \quad (x \in X).$$

Definition 2.10. Let A be a dual Banach algebra with identity. Then a net (M_α) in $L_{\omega^*}^2(A, C)^*$ is called an approximate normal, virtual diagonal for A if

$$a.M_\alpha - M_\alpha.a \longrightarrow 0 \quad \text{and} \quad \Delta_{\omega^*}(M_\alpha) \longrightarrow e$$

From now on we assume that A is a dual Banach algebra with identity. It is well known that every dual Banach algebra with a normal virtual diagonal is Connes-amenable [15]. In the following theorem we extend this result to approximate Connes-amenability.

Theorem 2.11. If A has an approximate normal, virtual diagonal $\{M_\alpha\}$, then A is approximately Connes-amenable.

Proof. Suppose X is a normal dual Banach A -bimodule with predual X_* and $D \in Z^1(A, X)$ $\omega^* - \omega^*$ -continuous. Since A is unital, by Proposition 3.11 we can assume that X is unital. We follow the argument in [4, Theorem 3.1]. The bilinear map $F(a, b) = Da.b$ is $\omega^* - \omega^*$ -continuous. Thus by the above remark we may let,

$$\phi_\alpha = \int F(a, b) dM_\alpha(a, b) = \int Da.bdM_\alpha \in X.$$

For $c \in A$, $x \in X_*$, we have

$$\begin{aligned} \langle c.\phi_\alpha, x \rangle &= \langle \phi_\alpha, x.c \rangle \\ &= \int \langle c.Da.b, x \rangle dM_\alpha(a, b) \\ &= \langle \int c.Da.b \ dM_\alpha(a, b), x \rangle. \end{aligned}$$

Therefore $c.\phi_\alpha = \int c.Da.b \ dM_\alpha(a, b)$ and similarly $\phi_\alpha.c = \int Da.bc \ dM_\alpha(a, b)$.

We have the following relations,

$$\int \langle D(ca).b, x \rangle dM_\alpha(a, b) = \langle c.M_\alpha, F_x \rangle \quad (1)$$

and

$$\int \langle Da.bc, x \rangle dM_\alpha(a, b) = \langle M_\alpha.c, F_x \rangle \quad (2)$$

where $F_x(a, b) = \langle Da.b, x \rangle$ and $F_x \in L_{\omega^*}^2(A, C)$.

By (1) and (2), we have

$$|\langle \int D(ca).b \ dM_\alpha(a, b) - \int Da.bc \ dM_\alpha(a, b), x \rangle| \leq \|c.M_\alpha - M_\alpha.c\| \|F_x\|$$

so

$$\| \int Dca.b dM_\alpha(a, b) - \int Da.bcdM_\alpha(a, b) \| \leq \|c.M_\alpha - M_\alpha.c\| \|D\| \|a\| \|b\| \quad (3).$$

On the other hand since $M_\alpha = \int a \otimes bdM_\alpha(a, b)$, then

$$\Delta_{\omega^*}(M_\alpha) = \int abdM_\alpha(a, b). \quad (4)$$

Thus

$$\begin{aligned} c.\phi_\alpha &= \int c.Da.b dM_\alpha(a, b) \\ &= \int D(ca).b dM_\alpha(a, b) - \int Dc.abdM_\alpha(a, b). \end{aligned}$$

So we have

$$\begin{aligned} c.\phi_\alpha - \phi_\alpha.c &= \int D(ca).b dM_\alpha(a, b) \\ &\quad - \int Da.bcdM_\alpha(a, b) - \int Dc.abdM_\alpha(a, b) \\ &= \int D(ca).b dM_\alpha(a, b) - \int Da.bcdM_\alpha(a, b) \\ &\quad - Dc. \int abdM_\alpha(a, b) \end{aligned}$$

By assumption and (3),(4) we have

$$Dc = \lim_{\alpha} \phi_\alpha.c - c.\phi_\alpha$$

$$= \lim_{\alpha} c.(-\phi_{\alpha}) - (-\phi_{\alpha}).c \quad (c \in A).$$

Thus D is approximately inner, and hence A is approximately Connes-amenable. $\square$

Definition 2.12. A is called approximately strongly Connes-amenable if for each unital Banach A -bimodule X , every $\omega^* - \omega^*$ continuous derivation $D : A \longrightarrow X^*$ whose range consists of $\omega^* -$ elements is approximately inner.

We don't know whether the converse of Theorem 2.11 is true, but for approximate strong Connes-amenable, the corresponding question is easy to answer:

Theorem 2.13. The following conditions are equivalent:

- (i) A has an approximate normal, virtual diagonal
- (ii) A is approximately strongly Connes-amenable.

Proof. (i) $\implies$ (ii). This is similar to Theorem 3.13.

(ii) $\implies$ (i) We follow the argument in [10, Theorem 4.7]. Since Δ_{ω^*} is $\omega^* - \omega^*$ continuous then $\ker \Delta_{\omega^*}$ is $\omega^* -$ closed and

$$(L_{\omega^*}^2(A, C)^*/{}^{\perp} \ker \Delta_{\omega^*})^* = \ker \Delta_{\omega^*}.$$

Clearly $ad_{e \otimes e}$ attains its values in the $\omega^* -$ elements of $\ker \Delta_{\omega^*}$. By the definition of approximate strong Connes-amenable, there exists a net $(N_{\alpha}) \subset \ker \Delta_{\omega^*}$ such that

$$ad_{e \otimes e}(a) = \lim_{\alpha} a.N_{\alpha} - N_{\alpha}.a \quad (a \in A).$$

Let $M_{\alpha} = e \otimes e - N_{\alpha}$. It follows that

$$a.M_{\alpha} - M_{\alpha}.a \longrightarrow 0 \quad \text{and} \quad \Delta_{\omega^*}(M_{\alpha}) \longrightarrow e \quad (a \in A). \quad \square$$

One of the unsatisfactory sides of dealing with approximate Connes-amenable for dual Banach algebras is the apparent lack of a suitable intrinsic characterization in terms of approximate normal, virtual diagonals. we saw that dual Banach algebras with an approximate normal, virtual diagonal are approximately Connes-amenable, but the converse is likely to be false in general. However for a compact group G the converse is true:

Theorem 2.14. Let G be a compact group. Then there is an approximate normal, virtual diagonal for $M(G)$.

Proof. By [12, Proposition 3.3] this is immediate. $\square$

We now modify the definition of approximate normal, virtual diagonal and obtain the desired characterization. Let A be a dual Banach algebra with predual A_* and let $\Delta : A \widehat{\otimes} A \longrightarrow A$ be the multiplication map. From [12, Corollary 4.6], we conclude that Δ^* maps A_* into $\sigma WC((A \widehat{\otimes} A)^*)$ [14]. Consequently, Δ^{**} induces the homomorphism

$$\Delta_{\sigma WC} : \sigma WC((A \widehat{\otimes} A)^*)^* \longrightarrow A.$$

With these preparations made, we can now characterize approximately Connes-amenable, dual Banach algebras through the existence of certain approximate normal, virtual diagonals. This is indeed an approximate version [12, Theorem 4.8].

Definition 2.15. An approximate σWC - virtual diagonal for A is a net $(M_\alpha) \subset \sigma WC((A \widehat{\otimes} A)^*)^*$ such that

$$a.M_\alpha - M_\alpha.a \longrightarrow 0 \quad \text{and} \quad \Delta_{\sigma WC}(M_\alpha) \longrightarrow e \quad (a \in A).$$

Theorem 2.16. The following conditions are equivalent:

- (i) A is approximately Connes-amenable.
- (ii) There is an approximate σWC - virtual diagonal for A .

Proof (i) $\implies$ (ii): First, note that $A \widehat{\otimes} A$ is canonically mapped into $\sigma WC((A \widehat{\otimes} A)^*)^*$. Define a derivation

$$D : A \longrightarrow \sigma WC((A \widehat{\otimes} A)^*)^*, \quad a \longmapsto a \otimes e - e \otimes a.$$

Since the dual module $\sigma WC((A \widehat{\otimes} A)^*)^*$ is normal, then it follows that D is ω^* -continuous.

Clearly, D attains its values in the ω^* -closed submodule $\text{Ker} \Delta_{\sigma WC}$. Since it is a normal dual Banach A -module, then there is a net $(N_\alpha) \subset \text{ker} \Delta_{\sigma WC}$ such that

$$Da = \lim_{\alpha} a.N_\alpha - N_\alpha.a \quad (a \in A).$$

Letting $M_\alpha = e \otimes e - N_\alpha$, we see that it is the approximate σWC -virtual diagonal for A .

(ii) $\implies$ (i) Let X be a normal, dual Banach A -bimodule. By Proposition 2.9 we may assume that X is unital, and let $D : A \longrightarrow X$ be a ω^* -derivation. Define

$$\theta_D : A \widehat{\otimes} A \longrightarrow X, \quad a \otimes b \longmapsto a.Db.$$

By [12, lemma 4.9], θ_D^* maps the predual X_* into $\sigma WC((A \widehat{\otimes} A)^*)$. Hence

$$(\theta^*|_{E_*})^* : \sigma WC((A \widehat{\otimes} A)^*)^* \longrightarrow E.$$

Let $(M_\alpha) \subset \sigma WC((A \widehat{\otimes} A)^*)^*$ be an approximate σWC -virtual diagonal for A , and let $x_\alpha = (\theta^*|_{E_*})^*(M_\alpha)$. We have the restriction map

$$\iota^* : (A \widehat{\otimes} A)^{**} \longrightarrow \sigma WC((A \widehat{\otimes} A)^*)^*.$$

Therefore we can choose a net $\sum_{k=1}^{n_\alpha} a_k^\alpha \otimes b_k^\alpha \in A \otimes A$ such that

$$M_\alpha = \omega^* - \lim_{\alpha} \sum_{k=1}^{n_\alpha} a_k^\alpha \otimes b_k^\alpha \quad \text{on} \quad \sigma WC((A \widehat{\otimes} A)^*)^*.$$

For every $c \in A$,

$$\begin{aligned} x_\alpha \cdot c &= \omega^* - \lim_{\alpha} \sum_{k=1}^{n_\alpha} a_k^\alpha \cdot D(b_k^\alpha) \cdot c \\ &= \omega^* - \lim_{\alpha} (\sum_{k=1}^{n_\alpha} a_k^\alpha \cdot D(b_k^\alpha c) - \sum_{k=1}^{n_\alpha} a_k^\alpha b_k^\alpha \cdot Dc) \\ &= (\theta^*|_{E_*})^*(M_\alpha \cdot c) - \Delta_{\sigma WC}(M_\alpha) \cdot Dc. \end{aligned}$$

Therefore

$$c \cdot x_\alpha - x_\alpha \cdot c = (\theta^*|_{E_*})^*(c \cdot M_\alpha - M_\alpha \cdot c) + \Delta_{\sigma WC}(M_\alpha) \cdot Dc. \quad (1)$$

So by assumption and (1), we have

$$Dc = \lim_{\alpha} c \cdot x_\alpha - x_\alpha \cdot c \quad (c \in A).$$

This implies that A is approximately Connes-amenable. $\square$

References

- [1] H. G. Dales, Banach algebras and automatic continuity, Clarendon Press, Oxford, 2000.
- [2] H. G. Dales, F. Gahramani, and N. Gronbeak, Derivations into iterated duals of Banach algebras, *studia mathematica* 128 (1) (1998), 19-53.
- [3] M. Daws, Connes-amenability of bidual and weighted semigroup algebras, *Math. Scand.* 99 (2006), no.2, 217-246.
- [4] E. D. Effros, Amenability and virtual diagonals for von Neumann algebras, *J. Funct. Anal.* 78 (1988), 137-153.
- [5] F. Ghahramani and R. J. Loy, Generalized notions of amenability, *J. Funct. Anal.* 208 (2004), 229-260.
- [6] F. Ghahramani, R. J. Loy and Y. Zhang, Generalized notions of amenability II, *J. Funct. Anal.* 254 (2008), 1776-1810.
- [7] A. Ya. Helemskii, Homological essence of amenability in the sense of A. Connes: the injectivity of the predual bimodule (translated from the Russian). *Math. USSR-Sb* 68 (1991), 555-566.
- [8] B. E. Johnson, Cohomology in Banach algebras, *Mem. Amer. Math. Soc.* 127 (1972).

- [9] B. E. Johnson, R. V. Kadison, and J. Ringrose, Cohomology of operator algebras, III. Bull. Soc. Math. France 100 (1972), 73-79.
- [10] V. Runde, Amenability for dual Banach algebras. Studia Math. 148 (2001), 47-66.
- [11] V. Runde, Lectures on amenability, Lecture Notes in Mathematics 1774, Springer-Verlage, Berlin, 2002.
- [12] V. Runde, Connes-amenability and normal virtual diagonals for measure algebras I, J. London Math. Soc. 67(2003), 643-656.
- [13] V. Runde, Connes-amenability and normal virtual diagonals for measure algebras II, Bull. Aust. Math. Soc. 68 (2003), 325-328.
- [14] V. Runde, Dual Banach algebras: Connes-amenability, normal, virtual diagonal, and injectivity of the predual bimodule (to appear).
- [15] E. G. Effros, Amenability and virtual diagonals for von Neumann algebras, J. Funct. Anal. 78 (1988), 137-156.

Acknowledgment

The authors would like to express their sincere thanks to Professor F. Ghahramani for his valuable comments.

G. H. Esslamzadeh

Department of Mathematics, Faculty of Sciences, Shiraz University, Shiraz 71454, Iran
e-mail: esslamz@shirazu.ac.ir

B. Shojaei

Islamic Azad University of Karadj, Karadj, Iran
e-mail: shoujaei@kiaui.ac.ir